

Generalized Weierstrass representation for surfaces in terms of Dirac-Hestenes spinor field

Vadim V. Varlamov

*Computer Division, Siberia State University of Industry,
Novokuznetsk 654007, Russia*

Abstract

A representation of generalized Weierstrass formulae for an immersion of generic surfaces into a 4-dimensional complex space in terms of spinors treated as minimal left ideals of Clifford algebras is proposed. The relation between integrable deformations of surfaces via mVN-hierarchy and integrable deformations of spinor fields on the surface is also discussed.

Mathematics Subject Classification (1991): 53A05, 53A10, 15A66.

Key words: Weierstrass representation, Clifford algebras, spinors.

1 Introduction

The theory of integrable deformations and immersions of surfaces due its a close relationship with the theory of integrable systems at present time is a rapid developing area of mathematical physics. One of the most powerful methods in this area is a Weierstrass representation for minimal surfaces [1], the generalization of which onto a case of generic surfaces was proposed by Konopelchenko in 1993 [2, 3] served as a basis for the following investigations. So, the generalized Weierstrass formulae for conformal immersion of surfaces into 3-dimensional Euclidean space are used for the study of the basic quantities related to 2D gravity, such as Polyakov extrinsic action, Nambu-Goto action, geometric action and Euler characteristic [4]. This method is also intensively used for the study of constant mean curvature surfaces, Willmore surfaces, surfaces of revolution and in many other problems related with differential geometry [5]-[14]. A further generalization of Weierstrass representation onto a case of multidimensional Riemann spaces, in particular onto a case of 4-dimensional space with signature $(+, +, +, -)$ (Minkowski space-time) has been proposed in the recent paper [15].

In the present paper we consider a relation between a Weierstrass representation in a 4-dimensional complex space $\mathbf{C}^4$ and a Dirac-Hestenes spinor field which is defined in Minkowski space-time $\mathbf{R}^{1,3}$. Dirac-Hestenes spinors were originally introduced in [16, 17] for the formulation of a Dirac theory of electron with the usage of the space-time algebra $\mathcal{C}_{1,3}$ [18] in $\mathbf{R}^{1,3}$ (see also [19]). On the other hand, there is a very graceful formulation [20]-[23] of the Dirac-Hestenes theory in terms of modern interpretation of spinors as minimal left ideals of Clifford algebras [24, 25], a brief review of which we give in Section 2. In Section 3 after a short historical introduction, generalized Weierstrass formulae in $\mathbf{C}^4$ are rewritten in a spinor representation type form (matrix representation of a biquaternion algebra $\mathbf{C}_2 \cong M_2(\mathbf{C})$) and are identified with the Dirac-Hestenes spinors, the matrix representation of which is also isomorphic to $M_2(\mathbf{C})$. It allows to use a well-known relation between Dirac-Hestenes and Dirac spinors [23, 26] (see also [27]) to establish a relation between Weierstrass-Konopelchenko coordinates for surfaces immersed into $\mathbf{C}^4$ and Dirac spinors. Integrable deformations of surfaces defined by a modified Veselov-Novikov equation and their relation with integrable deformations of Dirac field on surface are considered at the end of the Section 3.

2 Spinors as minimal left ideals of Clifford algebras

Let us consider a Clifford algebra $\mathcal{C}_{p,q}(V, Q)$ over a field $\mathbf{K}$ of characteristic 0 ($\mathbf{K} = \mathbf{R}$, $\mathbf{K} = \Omega = \mathbf{R} \oplus \mathbf{R}$, $\mathbf{K} = \mathbf{C}$), where V is a vector space endowed with a nondegenerate quadratic form

$$Q = x_1^2 + \dots + x_p^2 - \dots - x_{p+q}^2.$$

The algebra $\mathcal{C}_{p,q}$ is naturally $\mathbf{Z}_2$ -graded. Let $\mathcal{C}_{p,q}^+$ (resp. $\mathcal{C}_{p,q}^-$) be a set consisting of all even (resp. odd) elements of algebra $\mathcal{C}_{p,q}$. The set $\mathcal{C}_{p,q}^+$ is a subalgebra of $\mathcal{C}_{p,q}$. It is obvious that $\mathcal{C}_{p,q} = \mathcal{C}_{p,q}^+ \oplus \mathcal{C}_{p,q}^-$.

When n is odd, a volume element $\omega = \mathbf{e}_{12\dots p+q}$ commutes with all elements of algebra $\mathcal{C}_{p,q}$ and therefore belongs to a center of $\mathcal{C}_{p,q}$. Thus, in the case of n is odd we have for a center

$$\mathbf{Z}_{p,q} = \begin{cases} \mathbf{R} \oplus i\mathbf{R} & \text{if } \omega^2 = -1; \\ \mathbf{R} \oplus e\mathbf{R} & \text{if } \omega^2 = +1, \end{cases} \quad (1)$$

where e is a double unit. In the case of n is even the center of $\mathcal{C}_{p,q}$ consists the unit of algebra.

Let $\mathbf{R}_{p,q} = \mathcal{C}_{p,q}(\mathbf{R}^{p,q}, Q)$ be a real Clifford algebra ($V = \mathbf{R}^{p,q}$ is a real space). Analogously, in the case of a complex space we have $\mathbf{C}_{p,q} = \mathcal{C}_{p,q}(\mathbf{C}^{p,q}, Q)$. Moreover, it is obvious that $\mathbf{C}_{p,q} \cong \mathbf{C}_n$, where $n = p + q$. Further, let us consider the following most important in physics Clifford algebras and their isomorphisms to matrix algebras:

quaternions	$\mathbf{R}_{0,2} = \mathbb{H}$
biquaternions	$\mathbf{C}_2 = \mathbf{R}_{3,0} \cong \mathbf{M}_2(\mathbf{C})$
space-time algebra	$\mathbf{R}_{1,3} \cong \mathbf{M}_2(\mathbb{H})$
Dirac algebra	$\mathbf{C}_4 = \mathbf{R}_{4,1} \cong \mathbf{M}_4(\mathbf{C}) \cong \mathbf{M}_2(\mathbf{C}_2)$

The identity $\mathbf{C}_2 = \mathbf{R}_{3,0}$ for a biquaternion algebra known in physics as a Pauli algebra is immediately obtained from definition of the center of the algebra $\mathcal{C}_{p,q}$ (1). Namely, for $\mathbf{R}_{3,0}$ we have a volume element $\omega = \mathbf{e}_{123} \in \mathbf{Z}_{3,0} = \mathbf{R} \oplus i\mathbf{R}$, since $\omega^2 = -1$. The identity $\mathbf{C}_4 = \mathbf{R}_{4,1}$ is analogously proved. The isomorphism $\mathbf{R}_{4,1} \cong \mathbf{M}_2(\mathbf{C}_2)$ is a consequence of an algebraic modulo 2 periodicity of complex Clifford algebras: $\mathbf{C}_4 \cong \mathbf{C}_2 \otimes \mathbf{C}_2 \cong \mathbf{C}_2 \otimes \mathbf{M}_2(\mathbf{C}) \cong \mathbf{M}_2(\mathbf{C}_2)$ [28, 29, 30].

The left (resp. right) ideal of algebra $\mathcal{O}_{p,q}$ is defined by the expression $\mathcal{O}_{p,q}e$ (resp. $e\mathcal{O}_{p,q}$), where e is an idempotent satisfying the condition $e^2 = e$. Analogously, a minimal left (resp. right) ideal is a set of type $I_{p,q} = \mathcal{O}_{p,q}e_{pq}$ (resp. $e_{pq}\mathcal{O}_{p,q}$), where e_{pq} is a primitive idempotent, i.e., $e_{pq}^2 = e_{pq}$ and e_{pq} cannot be represented as a sum of two orthogonal idempotents, i.e., $e_{pq} \neq f_{pq} + g_{pq}$, where $f_{pq}g_{pq} = g_{pq}f_{pq} = 0$, $f_{pq}^2 = f_{pq}$, $g_{pq}^2 = g_{pq}$. In the general case a primitive idempotent has a form [20]

$$e_{pq} = \frac{1}{2}(1 + \mathbf{e}_{\alpha_1})\frac{1}{2}(1 + \mathbf{e}_{\alpha_2}) \dots \frac{1}{2}(1 + \mathbf{e}_{\alpha_k}), \quad (2)$$

where $\mathbf{e}_{\alpha_1}, \dots, \mathbf{e}_{\alpha_k}$ are commuting elements of the canonical basis of $\mathcal{O}_{p,q}$ such that $(\mathbf{e}_{\alpha_i})^2 = 1$, $(i = 1, 2, \dots, k)$. The values of k are defined by a formula

$$k = q - r_{q-p}, \quad (3)$$

where r_i are the Radon-Hurwitz numbers, values of which form a cycle of the period 8 :

$$r_{i+8} = r_i + 4. \quad (4)$$

The values of all r_i are

i	0	1	2	3	4	5	6	7
r_i	0	1	2	2	3	3	3	3

For example, let consider a minimal left ideal of the space-time algebra $\mathbf{R}_{1,3}$. The Radon-Hurwitz number for algebra $\mathbf{R}_{1,3}$ is equal to $r_{q-p} = r_2 = 2$, and therefore from (3) we have $k = 1$. The primitive idempotent of $\mathbf{R}_{1,3}$ has a form

$$e_{13} = \frac{1}{2}(1 + \mathbf{e}_0),$$

or $e_{13} = \frac{1}{2}(1 + \Gamma_0)$, where Γ_0 is a matrix representation of the unit $\mathbf{e}_0 \in \mathbf{R}_{1,3}$. Thus, a minimal left ideal of $\mathbf{R}_{1,3}$ is defined by the following expression

$$I_{1,3} = \mathbf{R}_{1,3}\frac{1}{2}(1 + \Gamma_0). \quad (5)$$

Analogously, for the Dirac algebra $\mathbf{R}_{4,1}$ on using the recurrence formula (4) we obtain $k = 1 - r_{-3} = 1 - (r_5 - 4) = 2$, and a primitive idempotent of $\mathbf{R}_{4,1}$ may be defined as follows

$$e_{41} = \frac{1}{2}(1 + \Gamma_0)\frac{1}{2}(1 + i\Gamma_{12}), \quad (6)$$

where $\Gamma_{12} = \Gamma_1\Gamma_2$ and Γ_i ($i = 0, 1, 2, 3$) are matrix representations of the units of $\mathbf{R}_{4,1} = \mathbf{C}_4$:

$$\begin{aligned}\Gamma_0 &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, & \Gamma_1 &= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}, \\ \Gamma_2 &= \begin{pmatrix} 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \\ 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \end{pmatrix}, & \Gamma_3 &= \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}.\end{aligned}$$

Further, for a minimal left ideal of Dirac algebra $I_{4,1} = \mathbf{R}_{4,1}\frac{1}{2}(1+\Gamma_0)\frac{1}{2}(1+i\Gamma_{12})$ using the isomorphisms $\mathbf{R}_{4,1} = \mathbf{C}_4 = \mathbf{C} \otimes \mathbf{R}_{1,3} \cong \text{M}_2(\mathbf{C}_2)$, $\mathbf{R}_{4,1}^+ \cong \mathbf{R}_{1,3} \cong \text{M}_2(\mathbb{H})$ and also an identity $\mathbf{R}_{1,3}e_{13} = \mathbf{R}_{1,3}^+e_{13}$ [22, 23] we have the following expression [27]:

$$I_{4,1} = \mathbf{R}_{4,1}e_{41} = (\mathbf{C} \otimes \mathbf{R}_{1,3})e_{41} \cong \mathbf{R}_{4,1}^+e_{41} \cong \mathbf{R}_{1,3}e_{41} = R_{1,3}e_{13}\frac{1}{2}(1+i\Gamma_{12}) = \mathbf{R}_{1,3}^+e_{13}\frac{1}{2}(1+i\Gamma_{12}). \quad (7)$$

Let $\Phi \in \mathbf{R}_{4,1} \cong \text{M}_4(\mathbf{C})$ be a Dirac spinor and $\phi \in \mathbf{R}_{1,3}^+ \cong \mathbf{R}_{3,0} = \mathbf{C}_2$ be a Dirac-Hestenes spinor. Then from (7) the relation immediately follows between spinors Φ and ϕ :

$$\Phi = \phi\frac{1}{2}(1+\Gamma_0)\frac{1}{2}(1+i\Gamma_{12}). \quad (8)$$

Since $\phi \in \mathbf{R}_{1,3}^+ \cong \mathbf{R}_{3,0}$, then the Dirac-Hestenes spinor can be represented by a biquaternion number

$$\phi = a^0 + a^{01}\Gamma_{01} + a^{02}\Gamma_{02} + a^{03}\Gamma_{03} + a^{12}\Gamma_{12} + a^{13}\Gamma_{13} + a^{23}\Gamma_{23} + a^{0123}\Gamma_{0123}. \quad (9)$$

Or in the matrix representation

$$\phi = \begin{pmatrix} \phi_1 & -\phi_2^* & \phi_3 & \phi_4^* \\ \phi_2 & \phi_1^* & \phi_4 & -\phi_3^* \\ \phi_3 & \phi_4^* & \phi_1 & -\phi_2^* \\ \phi_4 & -\phi_3^* & \phi_2 & \phi_1^* \end{pmatrix}, \quad \phi_i \in \mathbf{C}, \quad (10)$$

where

$$\phi_1 = a^0 - ia^{12}, \quad \phi_2 = a^{31} - ia^{23}, \quad \phi_3 = a^{03} - ia^{0123}, \quad \phi_4 = a^{01} + ia^{02}.$$

Finally, from (8) it follows that for the Dirac spinor Φ and also a space-time spinor $Z = \phi_2^1(1 + \Gamma_0)$ we have expressions

$$\Phi = \begin{pmatrix} \phi_1 & 0 & 0 & 0 \\ \phi_2 & 0 & 0 & 0 \\ \phi_3 & 0 & 0 & 0 \\ \phi_4 & 0 & 0 & 0 \end{pmatrix}, \quad Z = \begin{pmatrix} \phi_1 & -\phi_2^* & 0 & 0 \\ \phi_2 & \phi_1^* & 0 & 0 \\ \phi_3 & \phi_4^* & 0 & 0 \\ \phi_4 & -\phi_3^* & 0 & 0 \end{pmatrix},$$

which are minimal left ideals of algebras $\mathbf{R}_{4,1}$ and $\mathbf{R}_{1,3}$, respectively.

The Dirac spinor Φ may be considered as a vector in the 4-dimensional complex space $\mathbf{C}^4$ associated with the algebra $\mathbf{C}_4$. However, from a physical point of view it is more natural to consider the spinor Φ in space-time $\mathbf{R}^{1,3}$. In connection with this, let us introduce (follows [21, 22, 23, 26]) a more rigorous definition of spinor as a minimal left ideal of algebra $\mathcal{C}_{p,q}$.

Let $\mathfrak{B}_\Sigma = \{\Sigma_0, \dot{\Sigma}, \ddot{\Sigma}, \dots\}$ be a set of all ordered orthonormal bases for $\mathbf{R}^{p,q}$. Any two bases $\Sigma_0, \dot{\Sigma} \in \mathfrak{B}_\Sigma$ are related by the element of the group $\mathbf{Spin}_+(p, q)$:

$$\dot{\Sigma} = u\Sigma_0u^{-1}, \quad u \in \mathbf{Spin}_+(p, q).$$

Analogously, for the primitive idempotents defined in the basis $\Sigma \in \mathfrak{B}_\Sigma$ and denoted as $e_{\Sigma_0}, e_{\dot{\Sigma}}, \dots$, we have $e_{\dot{\Sigma}} = ue_{\Sigma_0}u^{-1}$, $u \in \mathbf{Spin}_+(p, q)$. Then the ideals $I_{\Sigma_0}, I_{\dot{\Sigma}}, I_{\ddot{\Sigma}}, \dots$ are geometrically equivalent if and only if

$$I_{\dot{\Sigma}} = uI_{\Sigma_0}u^{-1}, \quad u \in \mathbf{Spin}_+(p, q),$$

or, since $uI_{\Sigma_0} = I_{\Sigma_0}$:

$$I_{\dot{\Sigma}} = I_{\Sigma_0}u^{-1}.$$

Therefore, an algebraic spinor for $\mathbf{R}^{p,q}$ is an equivalence class of the quotient set $\{I_\Sigma\}/\mathbf{R}$, where $\{I_\Sigma\}$ is a set of all geometrically equivalent ideals, and $\Phi_{\Sigma_0} \in I_{\Sigma_0}$ and $\Phi_{\dot{\Sigma}} \in I_{\dot{\Sigma}}$ are equivalent, $\Phi_{\dot{\Sigma}} \cong \Phi_{\Sigma_0} \pmod{\mathbf{R}}$ if and only if $\Phi_{\dot{\Sigma}} = \Phi_{\Sigma_0}u^{-1}$, $u \in \mathbf{Spin}_+(p, q)$.

3 Weierstrass representation for surfaces in space $\mathbb{C}^4$

Historically, the Weierstrass representation [1] appeared in the result of the following variational problem: among the surfaces restricted by the some curve for finding such a surface, the area of which is minimal, i.e., it is necessary to find a minimum of the functional

$$S = \int \int \sqrt{1 + p^2 + q^2} dx dy,$$

where $p = dz/dx$, $q = dz/dy$, $z = f(x, y)$ is an equation of the surface. The Euler equation for this problem has a form

$$\frac{\partial}{\partial x} \left(\frac{p}{\sqrt{1 + p^2 + q^2}} \right) + \frac{\partial}{\partial y} \left(\frac{q}{\sqrt{1 + p^2 + q^2}} \right) = 0.$$

This equation expresses a main geometrical property of such a surface : in each point the meant curvature is equal to zero. The surface which possesses such a property is called *a minimal surface*. If we compare a region $\mathfrak{M}$ of the surface with a region $\mathfrak{E}$ of the flat surface so that the point on $\mathfrak{M}$ with the coordinates (X^1, X^2, X^3) corresponds to a point $w = u + iv$ of region $\mathfrak{E}$, then for the minimal surface we have the equations

$$\frac{\partial^2 X^1}{\partial u^2} + \frac{\partial^2 X^1}{\partial v^2} = 0, \quad \frac{\partial^2 X^2}{\partial u^2} + \frac{\partial^2 X^2}{\partial v^2} = 0, \quad \frac{\partial^2 X^3}{\partial u^2} + \frac{\partial^2 X^3}{\partial v^2} = 0,$$

solutions of which are of the form

$$X^1 = \operatorname{Re} f(w), \quad X^2 = \operatorname{Re} g(w), \quad X^3 = \operatorname{Re} h(w),$$

at

$$(f'(w))^2 + (g'(w))^2 + (h'(w))^2 = 0.$$

The functions satisfying this equation are

$$f'(w) = i(G^2 + H^2), \quad g'(w) = G^2 - H^2, \quad h'(w) = 2GH,$$

whence

$$\begin{aligned}
X^1 &= C^1 + \operatorname{Re} \int_{w_0}^w i(G^2 + H^2)dw, \\
X^2 &= C^2 + \operatorname{Re} \int_{w_0}^w (G^2 - H^2)dw, \\
X^3 &= C^3 + 2\operatorname{Re} \int_{w_0}^w GHdw.
\end{aligned} \tag{11}$$

Here $G(w)$ and $H(w)$ are holomorphic functions defined in a circle or in all complex plane. After substitution of variables

$$s = \xi + i\eta = \frac{H(w)}{G(w)}, \quad G^2 \frac{dw}{ds} = F(s),$$

the equations (11) take the form

$$\begin{aligned}
dX^1 &= \operatorname{Re} [i(1 + s^2)F(s)ds], \\
dX^2 &= \operatorname{Re} [(1 - s^2)F(s)ds], \\
dX^3 &= \operatorname{Re} [2sF(s)ds].
\end{aligned}$$

Thus, for an every analytic function $F(s)$ we have a minimal surface.

Further, let us consider generalized Weierstrass representation for surfaces immersed into 4-dimensional complex space $\mathbf{C}^4$, which, as known, is associated with the Dirac algebra $\mathbf{C}_4$. In this case generalized Weierstrass formulae have a form

$$\begin{aligned}
X^1 &= \frac{i}{2} \int_{\Gamma} (\psi_1 \psi_2 d\bar{z} - \varphi_1 \varphi_2 dz), \\
X^2 &= \frac{1}{2} \int_{\Gamma} (\psi_1 \psi_2 d\bar{z} + \varphi_1 \varphi_2 dz), \\
X^3 &= \frac{1}{2} \int_{\Gamma} (\psi_1 \varphi_2 d\bar{z} - \varphi_1 \psi_2 dz), \\
X^4 &= \frac{i}{2} \int_{\Gamma} (\psi_1 \varphi_2 d\bar{z} + \varphi_1 \psi_2 dz),
\end{aligned} \tag{12}$$

$$\begin{aligned}
\psi_{\alpha z} &= p\varphi_{\alpha}, \\
\varphi_{\alpha \bar{z}} &= -p\psi_{\alpha}
\end{aligned}, \quad \alpha = 1, 2, \tag{13}$$

where ψ, φ and p are complex-valued functions on variables $z, \bar{z} \in \mathbf{C}$, Γ is a contour in complex plane $\mathbf{C}$. We will interpret the functions $X^i(z, \bar{z})$ as the coordinates in $\mathbf{C}^4$. It is easy to verify that components of an induced metric have a form

$$\begin{aligned} g_{zz} &= \overline{g_{\bar{z}\bar{z}}} = \sum_{i=1}^4 (X_z^i)^2 = 0, \\ g_{z\bar{z}} &= \sum_{i=1}^4 (X_z^i X_{\bar{z}}^i) = \psi_1 \psi_2 \varphi_1 \varphi_2. \end{aligned}$$

Therefore, the formulae (12), (13) define a conformal immersion of the surface into $\mathbf{C}^4$ with an induced metric

$$ds^2 = \psi_1 \psi_2 \varphi_1 \varphi_2 dz d\bar{z}.$$

The formulae (12) may be rewritten in the following form

$$\begin{aligned} d(X^1 + iX^2) &= i\psi_1 \psi_2 \bar{z}, \\ d(X^1 - iX^2) &= -i\varphi_1 \varphi_2 dz, \\ d(X^4 + iX^3) &= i\psi_1 \varphi_2 \bar{z}, \\ d(X^4 - iX^3) &= i\varphi_1 \psi_2 dz, \end{aligned}$$

or

$$d(X^4 \sigma_0 + X^1 \sigma_1 + X^2 \sigma_2 + X^3 \sigma_3) = i \begin{pmatrix} \varphi_1 \psi_2 dz & \psi_1 \psi_2 d\bar{z} \\ \varphi_1 \varphi_2 dz & \psi_1 \varphi_2 d\bar{z} \end{pmatrix}, \quad (14)$$

where

$$\sigma_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}$$

are matrix representations of the units of quaternion algebra $\mathbf{R}_{0,2} = \mathbb{H} : \mathbf{e}_i \longrightarrow \sigma_i$ ($i = 0, 1, 2$), $\mathbf{e}_{21} \longrightarrow \sigma_3$. It is easy to see that the left part of the expression (14) is a biquaternion $\mathbf{C}_2 = \mathbf{C} \otimes \mathbf{R}_{0,2}$. Recalling that $\mathbf{C}_2 = \mathbf{R}_{3,0}$ and a volume element $\omega = \mathbf{e}_{123} \in \mathbf{R}_{3,0}$ belongs to a center $\mathbf{Z}_{3,0} = \mathbf{R} \oplus i\mathbf{R}$, we can write a biquaternion $X^4 \mathbf{e}_0 + X^1 \mathbf{e}_1 + X^2 \mathbf{e}_2 + X^3 \mathbf{e}_3$,

where $\mathbf{e}_1^2 = \mathbf{e}_2^2 = 1$, $\mathbf{e}_3 = \mathbf{e}_{21} = \mathbf{e}_2\mathbf{e}_1$, in the form

$$\begin{aligned} & \text{Re } X^4 \mathbf{e}_0 + \text{Re } X^1 \mathbf{e}_1 + \text{Re } X^2 \mathbf{e}_2 + \text{Re } X^3 \mathbf{e}_3 + \\ & + \text{Im } X^3 \mathbf{e}_{12} + \text{Im } X^2 \mathbf{e}_{31} + \text{Im } X^1 \mathbf{e}_{23} + \text{Im } X^4 \mathbf{e}_{123} = \\ & = (\text{Re } X^4 + \omega \text{Im } X^4) \mathbf{e}_0 + (\text{Re } X^1 + \omega \text{Im } X^1) \mathbf{e}_1 + \\ & + (\text{Re } X^2 + \omega \text{Im } X^2) \mathbf{e}_2 + (\text{Re } X^3 + \omega \text{Im } X^3) \mathbf{e}_3 = \\ & = X^4 \mathbf{e}_0 + X^1 \mathbf{e}_1 + X^2 \mathbf{e}_2 + X^3 \mathbf{e}_3. \end{aligned} \quad (15)$$

Further, by means of isomorphisms $\mathbf{R}_{3,0} \cong \mathbf{R}_{1,3}^+$ and $\mathbf{R}_{4,1}^{++} \cong \mathbf{R}_{1,3}^+ \cong \mathbf{R}_{3,0}$ the biquaternion (15) may be rewritten as (like (9)):

$$\begin{aligned} \phi = & \text{Re } X^4 I + \text{Re } X^1 \Gamma_{01} + \text{Re } X^2 \Gamma_{02} + \text{Re } X^3 \Gamma_{03} + \\ & + \text{Im } X^3 \Gamma_{12} + \text{Im } X^2 \Gamma_{31} + \text{Im } X^1 \Gamma_{23} + \text{Im } X^4 \Gamma_{0123}. \end{aligned} \quad (16)$$

Or in the form (10) if suppose

$$\begin{aligned} \phi_1 &= \text{Re } X^4 - i \text{Im } X^3, \\ \phi_2 &= \text{Im } X^2 - i \text{Im } X^1, \\ \phi_3 &= \text{Re } X^3 - i \text{Im } X^4, \\ \phi_4 &= \text{Re } X^1 + i \text{Re } X^2. \end{aligned} \quad (17)$$

The formulae (17) define a relation between Weierstrass-Konopelchenko coordinates and Dirac-Hestenes spinors. This relation is a direct consequence of an isomorphism $\mathbf{C}_2 = \mathbf{C} \otimes \mathbf{R}_{0,2} = \mathbf{R}_{3,0} \cong \mathbf{R}_{1,3}^+$. Further, using the idempotent $\frac{1}{2}(1 + \Gamma_0)\frac{1}{2}(1 + i\Gamma_{12})$ it is easy to establish (by means of (8)) a relation with the Dirac spinor treated as a minimal left ideal of algebra $\mathbf{R}_{4,1} = \mathbf{C}_4 \cong \mathbf{M}_4(\mathbf{C})$:

$$\Phi = \begin{pmatrix} \phi_1 & 0 & 0 & 0 \\ \phi_2 & 0 & 0 & 0 \\ \phi_3 & 0 & 0 & 0 \\ \phi_4 & 0 & 0 & 0 \end{pmatrix}$$

It is obvious that we cannot directly to identify the spinor defined by the formulae (17) with a generic “physical” spinor of electron theory, because in accordance with (17) and (12)-(13) the spinor ϕ depends only on two variables $z, \bar{z}$, or x^1, x^2 if suppose $z = x^1 + ix^2$, whilst a physical spinor with four components depends on four variables x^1, x^2, x^3, x^4 . By this reason we will call the spinor defined by the identities (17) as *a surface spinor*,

and respectively the field $\Phi = (\phi_1, \phi_2, \phi_3, \phi_4)^T$ will be called a *Dirac spinor field on surface*. The relationship between a surface spinor $\phi(z, \bar{z})$ and a physical (space) spinor $\phi(x^1, x^2, x^3, x^4)$, and also a relation with the spinor representations of surfaces in spaces $\mathbf{R}^{p,q}$ will be considered in a separate paper.

Further, according to Section 2, an algebraic Dirac spinor for $\mathbf{R}^{1,3}$ is an element of $\{I_\Sigma\}/\mathbf{R}$. Then if $\Phi_{\Sigma_0} \in I_{\Sigma_0}$, $\Phi_{\dot{\Sigma}} \in I_{\dot{\Sigma}}$, then $\Phi_{\dot{\Sigma}} \simeq \Phi_{\Sigma_0} \pmod{\mathbf{R}}$ if and only if

$$\Phi_{\dot{\Sigma}} = \Phi_{\Sigma_0} u^{-1}, \quad u \in \mathbf{Spin}_+(1, 3). \quad (18)$$

Here in accordance with (8)

$$\Phi_{\Sigma_0} = \phi_{\Sigma_0} \frac{1}{2}(1 + \Gamma_0) \frac{1}{2}(1 + i\Gamma_{12}).$$

The formula (18) defines a transformation law of the Dirac spinor. It is obvious that a transformation group of the biquaternion (14) is also isomorphic to $\mathbf{Spin}_+(1, 3)$, since

$$\mathbf{Spin}_+(1, 3) \cong \left\{ \begin{pmatrix} a & c \\ b & d \end{pmatrix} \in \mathbf{C}_2 : \det \begin{pmatrix} a & c \\ b & d \end{pmatrix} = 1 \right\} = SL(2; \mathbf{C}),$$

where $SL(2; \mathbf{C})$ is a double covering of the own Lorentz group $\mathcal{L}_+^\uparrow$. Therefore, the transformations of Weierstrass-Konopelchenko coordinates for surfaces immersed into $\mathbf{C}^4$ are induced (via the relations (17)) transformation of a Dirac field $\Phi = (\phi_1, \phi_2, \phi_3, \phi_4)^T$ in $\mathbf{R}^{1,3}$, where $\Phi \in M_4(\mathbf{C})e_{41}$ is the minimal left ideal of $\mathbf{R}_{4,1} \cong M_4(\mathbf{C})$ defined in some orthonormal basis $\Sigma \in \mathfrak{B}_\Sigma$.

On the other hand, if suppose (following [2, 3, 15]) that the functions p, ψ_α and φ_α in (13) depend on the deformation parameter t , then the deformations of ψ_α and φ_α are defined by a following system:

$$\begin{aligned} \psi_{\alpha t} &= A\psi_\alpha + B\varphi_\alpha, \\ \varphi_{\alpha t} &= C\psi_\alpha + D\varphi_\alpha, \end{aligned} \quad \alpha = 1, 2 \quad (19)$$

where A, B, C, D are differential operators. The equations (19) define integrable deformations of surfaces immersed in $\mathbf{C}^4$. Let p be a real-valued function; the compatibility condition of (19) with (13) is equivalent to the nonlinear partial differential equation for p . In the simplest nontrivial case

(A, B, C, D are first order operators) it is a modified Veselov-Novikov equation [31]:

$$p_t + p_{zzz} + p_{\bar{z}\bar{z}\bar{z}} + 3p_z\omega + 3p_{\bar{z}}\bar{\omega} + \frac{3}{2}p\bar{\omega}_{\bar{z}} + \frac{3}{2}p\omega_z = 0,$$

$$\omega_{\bar{z}} = (p^2)_z.$$

Varying operators A, B, C, D one gets an infinite hierarchy of integrable equations for p (modified Veselov-Novikov hierarchy [32, 33, 31, 2]). It is obvious that the deformation of $\psi_\alpha, \varphi_\alpha$ via (19) induced the deformations of the coordinates $X^i(z, \bar{z}, t)$ in $\mathbf{C}^4$. Moreover, according to (14) and (16) treated as a matrix representation of the Dirac-Hestenes spinor field ϕ , we may say that the mVN-deformation generates a deformation of the Dirac field $\Phi = (\phi_1, \phi_2, \phi_3, \phi_4)^T$.

Acknowledgements

I am grateful to Prof. B.G. Konopelchenko whose articles and preprints are the ground for this work.

References

- [1] K. Weierstrass, *Untersuchungen über die Flächen, deren mittlere Krümmung überall gleich Null ist*, Monatsber. Akad. Wiss. Berlin, 1866, S. 612-625.
- [2] B.G. Konopelchenko, *Induced surfaces and their integrable dynamics*, Stud. Appl. Math., **96**, 9-51, (1996); preprint Institute of Nuclear Physics, N 93-144, Novosibirsk, (1993).
- [3] B.G. Konopelchenko, *Multidimensional integrable systems and dynamics of surfaces in space*, preprint of Institute of Mathematics, Taipei, August, 1993.
- [4] R. Carroll and B.G. Konopelchenko, *Generalized Weierstrass- Enneper inducing, conformal immersion and gravity*, Int. J. Modern Physics **A11**, 1183-1216, (1996).

- [5] B.G. Konopelchenko and I.A. Taimanov, *Generalized Weierstrass formulae, soliton equations and Willmore surfaces*, preprint N.187, Univ. Bochum, (1995).
- [6] B.G. Konopelchenko and I.A. Taimanov, *Constant mean curvature surfaces via an integrable dynamical system*, J. Phys. A: Math. Gen., **29**, 1261-1265, (1996).
- [7] I.A. Taimanov, *Modified Novikov-Veselov equation and differential geometry of surfaces*, Trans. Amer. Math. Soc., Ser.2, **179**, 133-159, (1997).
- [8] I.A. Taimanov, *Surfaces of revolution in terms of solitons*, Annals of Global Analysis and Geometry, **15**, N 5, 419-435, (1997).
- [9] I.A. Taimanov, *The Weierstrass representation of closed surfaces in R^3* , preprint dg-ga/9710020, 1997.
- [10] I.A. Taimanov, *Global Weierstrass representation and its spectrum*, Uspechi Mat. Nauk, **52**, N 6, 187-188, (1997).
- [11] P.G. Grinevich and M.V. Schmidt, *Conformal invariant functionals of immersion of tori into R^3* , J. Geometry and Physics **26**,1-2, 51-78, (1998).
- [12] B.G. Konopelchenko and G. Landolfi, *On classical string configurations*, Mod. Phys. Lett. A, **12**, (1997), 3161-3168.
- [13] R. Kusner and N. Schmitt, *The spinor representation of surfaces in space*, preprint dg-ga/9610005, 1996.
- [14] T. Friedrich, *On the spinor representation of surfaces in euclidean 3-spaces*, preprint SFB 288, N 295, TU-Berlin, 1997; J. Geometry and Physics (to appear).
- [15] B.G. Konopelchenko and G. Landolfi, *Generalized Weierstrass representation for surfaces in multidimensional Riemann spaces*, preprint math.DG/9804144, 1998; J. Geometry and Physics (to appear).
- [16] D. Hestenes, *Real spinor fields*, J. Math. Phys. **8**, 798-808, (1967).
- [17] D. Hestenes, *Observables, operators, and complex numbers in the Dirac theory*, J. Math. Phys. **16**, 556-571, (1976).

- [18] D. Hestenes, *Space-Time Algebra*, Gordon and Breach, New York, 1987.
- [19] D. Hestenes and G. Sobczyk, *Clifford algebra to Geometrical Calculus*, D. Heidel, Publ. Co. Dordrecht, 1984.
- [20] P. Lounesto, *Scalar product of spinors and an extension of Brauer-Wall groups*, Found. Phys. **11**, 721-740, (1981).
- [21] P. Lounesto, *Clifford algebras and Hestenes spinors*, Found. Phys. **23**, 1203-1237, (1993).
- [22] V.L. Figueiredo, W.A. Rodrigues, Jr. and E.C. Oliveira, *Covariant, algebraic, and operator spinors*, Int. J. Theor. Phys. **29**, 371-395, (1990).
- [23] V.L. Figueiredo, W.A. Rodrigues, Jr. and E.C. Oliveira, *Clifford algebras and the hidden geometrical nature of spinors*, Algebras, Groups and Geometries, **7**, 153-198, (1990).
- [24] A. Crumeyrolle, *The primitive idempotents of the Clifford algebras and the amorphic spinor fibre bundles*, Reports on Math. Phys. **25**, 305-328, (1987).
- [25] A. Crumeyrolle, *Orthogonal and Symplectic Clifford Algebras*, Kluwer Acad. Publ., Dordrecht, 1991.
- [26] W.A. Rodrigues, Jr., Q.A.G. De Souza, J. Vaz, Jr., and P. Lounesto, *Dirac-Hestenes spinor fields in Riemann-Cartan spacetime*, Int. J. Theor. Phys. **35**, 1849-1900, (1996).
- [27] J. Vaz, Jr., and W.A. Rodrigues, Jr., Maxwell and Dirac theories as an already unified theory, preprint hep-th/9511181, to appear in the proceedings of the International Conference on the Theory of the Electron, J. Keller and Oziewicz (eds.), UNAM, Mexico, (1995).
- [28] M.F. Atiyah, R. Bott and A. Shapiro, *Clifford modules*, Topology, **3**, (Suppl.1), 3-38, (1964).
- [29] I. Porteous, *Topological Geometry*, van Nostrand, London, 1969.
- [30] M. Karoubi, *K-Theory*, Springer-Verlag, Berlin, 1979.

- [31] L.V. Bogdanov, *Veselov-Novikov equation as a natural two-dimensional generalization of the Korteweg-de Vries equation*, Theor. Math. Phys. **70**, 309-314, (1987).
- [32] A.P. Veselov and S.P. Novikov, *Finite-zone, two-dimensional potential Schrödinger operators. Explicit formulas and evolution equations*, Soviet Math. Dokl. **30**, 588-591, (1984).
- [33] A.P. Veselov and S.P. Novikov, *Finite-zone, two-dimensional Schrödinger operators. Potential operators*, Soviet Math. Dokl. **30**, 705-708, (1984).